
(*Circus aeruginosus*)

(// : // :)

*

Archive of SID

/

:

.()

()

.()

o ')
o ')
(o ')

(Ss Ts,R, P, W)

()

(1a, 2a, 3a, 3c) 1, 2, 5, 6

.()

(*Anas strepera*)

(*Aythya fuligula*)

(*Fulica atra*)

()

.()

(*Pelecanus crispus*)

(*Anser erythropus*)

(*Aquila heliaca*)

(*Oxyura leucocephala*)

()

()

()

()

)
(

()

(*Circus aeruginosus*)

(Accipitridae)

(Aves)

(Falconiformes)

()

)

(

)

(

)

(Dr)

(

(Wdf)

(Wds)

(Graz)

...

) (()
 ()
 ()
) (Dam) ()
 ()
 (Wuse) ()
 () ()
 ()

Archive of SID

() Smith Hill

Smith Hill
 ADE-4 ()

Smith Hill

Total Inertia= (Inertia MCOA × Nb Var. Qualitative / NBVar.Tot.) + (Inertia PCA / NBVar.Tot.)

Inertia

Standardized Principal Component Analysis
 Ordination
 3 Multiple Correspondence Analysis

:

Smith Hill

1	+ ,625	+0.469	+0.469
2	+ ,625	+0.192	+0.661
3	+ ,625	+0.140	+0.802

() R² R²
:

Smith Hill

R ²				


() Smith Hill
(Dr)
(Wds Wdf)
()

(Wuse)
)

(Dam


(Am1-Am3)

(Graz)



.Smith Hill

Al = Aj = Am = ()
 .Si = ()



.Smith Hill

()

(A17-A117)

()

Wdf)

(Dam

(Dr)

)

(

(Si1-Si6)

)

(

)

(Dr Wdf

Aj1)

()

(Aj2

(Aj3 – Aj11)

)

(Graz

(

)

Archive of SID

()

...

P.)

(*Podiceps cristatus*)

(*Fulica atra*) (*ruficollis*

... (*Gallinula chloropus*)

(*Vulpes vulpes*)

(*Canis aureus*) (*V. corsac*)

(*Felis catus*)

()

Archive of SID



...
)

(

Archive of SID

5– Andelman, S. J., W. F. Fagen. 2000. Umbrella and flagships: efficient conservation surrogates or expensive mistakes. *Proceeding of the National Academy of Science* 97:5954-5959.

6– Berger, J. 1997. Population constraints associated with the use of black rhinos as an umbrella species for desert herbivores. *Conservation Biology* 11: 69-78.

7– Cramp S (eds) 1988. *Handbook of the Birds of Europe, the Middle East and North Africa, The Birds of Western Palaeartic, Vol. 5*. Oxford University Press, Oxford.

-
- 8– Denormandie, J., T. C. Edwards. 2001. The Umbrella concept and regional conservation planning in southern California: a comparative study. In review, *Conservation Biology*. April 2001.
- 9– Dickinson, E. C. 2003. *The Howard and Moore Complete Checklist of the Birds of the World*. London, Christopher Helm.
- 10– Evans, M.I. 1994. *Important Bird Areas in the Middle East*, Birdlife International Inc. 410 p..
- 11– Jongman, R. H., ter Braak, C.J.F. & van Tongeren, O.F.R. 1987. *Data analysis in community and landscape ecology*. Pudoc, Wageningen. 298 p..
- 12– Pitelka, F. A. 1981. The condor case: an uphill struggle in a downhill crush. *Auk* 98:634-635.
- 13– Sanjayan, M. A., Richard M. Jeo and Dennis Sizemore. 2003. Key Concept, Umbrella species. [On line] (Consultation 12/15/2003). Address URL. [WWW.savethegreatbear.org/CAO/Key concepts.htm](http://WWW.savethegreatbear.org/CAO/Key%20concepts.htm).
- 14– Sibley, C. G. and B. L. Monroe 1990. *Distribution and Taxonomy of Birds of the world*. London, Yale University Press.
- 15– Thioulouse J, Chessel D, Dolédec S, Olivier JM 1997. *ADE - 4 : Ecological data analysis: Exploratory and Eucliden methods in Environmental Sciences*, University of Lyon 1, Lyon, France.
- 16– Wetlands International, Ramsar Sites Database Service. [On line] (Consultation 01/10/2004). Address URL: <http://www.wetlands.org/RSDB/default.htm>.
- 17– Wilcox, B. A. 1984. In situ conservation of genetic resources: determinants of minimum area requirements. Pages 639-647 in J. A. McNeely, K. R. Miller, editors, *National Parks, conservation, and development: the role of protected areas in sustaining society*. Smithsonian, Institution Press, Washington D. C., USA.

A Study of the Factors affecting the Breeding Success of Marsh Harrier (*Circus aeruginosus*) as an Umbrella Species in Aji-gol, Ala-gol and Alma-gol International Wetlands of Turkmen Steppe

M.Kaboli^{*1}, M.Karami² and B.Kiabi³

¹Asst. Prof., Faculty of Natural Resources, University of Tehran, I. R. Iran

² Prof., Faculty of Natural Resources, University of Tehran, I. R. Iran

³ Assoc. Prof., Zoology Dep. of Beheshti University, Tehran, I. R. Iran

(Received: 22 Nov 2000, Accepted: 26 Sep 2005)

Abstract

This study was conducted to evaluate the problems and threats of Aji-gol, Ala-gol and Alma-gol international wetlands of Turkmen steppe in Golestan province, taking the Marsh harrier (*Circus aeruginosus*) as an umbrella species. Monitored were 31 nests, 131 eggs, 62 nestlings and 28 juveniles during two breeding periods, from late winter 1997 to late spring 1999. Factors affecting the nests and the nestlings were also regularly monitored. The three areas considered together, the breeding success of Marsh harrier was estimated to be about 21.37 %, a proportion much lower than that normally observed in Europe. Flooding of breeding habitats through construction of dams in the north-west of Ala-gol, and on the contrary, drying up of these habitats through drainage for aquaculture activities in the northern part and as well impact of livestock grazing are responsible for this breeding failure and consequently, the most important threats for the wetlands and their wildlife. In total, the current overexploitation of the areas with no due consideration of the seasonal needs of wildlife represents the principle threat to the biodiversity of the wetlands.

Keywords: International wetlands, Aji-gol, Ala-gol, Alma-gol, Marsh harrier, Umbrella species, Breeding success, Threats.

* Corresponding author: Tel:

, Fax:

Email: mkaboli@yahoo.com